

An atomic structure of human γ -secretase

Xiao-chen Bai^{1*}, Chuangye Yan^{2*}, Guanghui Yang^{2*}, Peilong Lu², Dan Ma², Linfeng Sun², Rui Zhou², Sjors H. W. Scheres¹ & Yigong Shi²

Dysfunction of the intramembrane protease γ -secretase is thought to cause Alzheimer's disease, with most mutations derived from Alzheimer's disease mapping to the catalytic subunit presenilin 1 (PS1). Here we report an atomic structure of human γ -secretase at 3.4 Å resolution, determined by single-particle cryo-electron microscopy. Mutations derived from Alzheimer's disease affect residues at two hotspots in PS1, each located at the centre of a distinct four transmembrane segment (TM) bundle. TM2 and, to a lesser extent, TM6 exhibit considerable flexibility, yielding a plastic active site and adaptable surrounding elements. The active site of PS1 is accessible from the convex side of the TM horseshoe, suggesting considerable conformational changes in nicastrin extracellular domain after substrate recruitment. Component protein APH-1 serves as a scaffold, anchoring the lone transmembrane helix from nicastrin and supporting the flexible conformation of PS1. Ordered phospholipids stabilize the complex inside the membrane. Our structure serves as a molecular basis for mechanistic understanding of γ -secretase function.

A hallmark of Alzheimer's disease is accumulation of β -amyloid plaque in the brains of patients¹. The intramembrane protease γ -secretase is thought to contribute to the development of Alzheimer's disease by generating β -amyloid peptides (A β s), particularly those that are prone to aggregation such as A β 42 (refs 2–5). A mature γ -secretase consists of four components: presenilin, PEN-2, nicastrin, and APH-1 (ref. 6). Among these components, presenilin is responsible for the A β -producing proteolytic activity^{7,8}.

Presenilin comprises nine TMs, with the signature motifs YD on TM6 and GxGD on TM7 (ref. 8). During assembly of γ -secretase, presenilin undergoes an autocatalytic cleavage to produce an amino (N)-terminal fragment (NTF, comprising TMs 1–6) and a carboxy (C)-terminal fragment (CTF, comprising TMs 7–9)⁹. Among the 300 or so mutations derived from patients with familial Alzheimer's disease (FAD), more than two-thirds are mapped to PS1, and about three dozen each are derived from PS2 and amyloid precursor protein (APP). Nicastrin contains a large extracellular domain (ECD) and a single TM; its ECD is heavily glycosylated and thought to recognize the N terminus of substrate protein^{10–13}. PEN-2 directly binds PS1 and is required for its autocatalytic maturation and protease activity^{14,15}. APH-1 contains seven TMs and is indispensable for γ -secretase assembly^{16,17}.

Our previous cryo-electron microscopy (cryo-EM) structure of human γ -secretase at 4.5 Å resolution led to identification of 19 TMs and construction of a partial atomic model for the ECD¹⁸. Analysis of the crystal structures of the archaeal presenilin homologue PSH¹⁹ and the nicastrin homologue from *Dictyostelium purpureum* (DpNCT)²⁰ yielded a tantalizing clue on TM assignment and an improved atomic model for the ECD. A subsequent cryo-EM structure of human γ -secretase at 4.3 Å resolution allowed assignment of all 20 TMs to the four components²¹. In this study, we report the first atomic structure of an intact human γ -secretase, which allows visualization of the atomic structures of all four components and determination of the specific interactions that underlie γ -secretase assembly.

Structure determination of human γ -secretase

We performed cryo-EM single-particle analysis on the same sample in amphipols that was used to calculate our previous map¹⁸. With the aim of reaching higher resolution, we used zero-loss energy-filtered imaging, a higher magnification, and a lower dose rate on the single-electron counting detector (see Methods). We also collected a larger data set. An initial set of 412,272 particle images yielded a 3.5 Å map. Subsequent three-dimensional (3D) classification that was focused on the TMs resulted in a more structurally homogeneous subset of 159,549 particles. This subset was used to calculate the final map at 3.4 Å resolution (Extended Data Figs 1 and 2). Our map displays excellent main-chain connectivity and side-chain densities for almost all residues (Extended Data Fig. 2a).

Twenty TMs were identified in the transmembrane region, including a highly mobile TM (TM2 of PS1) that is only visible as rod-shaped density in a 7 Å low-pass filtered map (Extended Data Fig. 2b). This low-pass filtered map also shows a second rod-shaped density in the PS1 cavity between TM3, TM5, and TM2, which we were unable to identify. As previously reported^{20,21}, PS1 and APH-1 are located at the centre of the TM horseshoe. All seven TMs of APH-1 have exceptional density, with most side chains clearly identifiable (Extended Data Fig. 2c). Except for TM2 and TM6, all other TMs of PS1 exhibit excellent density, with aromatic and bulky residues easily recognizable (Extended Data Fig. 2d). The three TMs of PEN-2 on the thin end of the TM horseshoe and the lone TM of nicastrin on the thick end both display discernible side-chain features (Extended Data Fig. 2e, f). Several N-linked glycans and lipid molecules are also defined by clear EM density (Extended Data Fig. 2g, h).

On the basis of these unambiguous EM densities, we built and refined a near-complete atomic model for human γ -secretase (Fig. 1 and Extended Data Table 1), which includes 598 residues in the TMs and 632 residues in the ECD. Assignment of specific residues in the TMs was aided by a large number of aromatic amino acids (Extended Data Fig. 2c–f). The density map for PS1-TM2 is inadequate for

¹MRC Laboratory of Molecular Biology, Cambridge Biomedical Campus, Cambridge CB2 0QH, UK. ²Ministry of Education Key Laboratory of Protein Science, Tsinghua-Peking Joint Center for Life Sciences, Center for Structural Biology, School of Life Sciences, Tsinghua University, Beijing 100084, China.

*These authors contributed equally to this work.




Figure 1 | Atomic structure of human γ -secretase. **a**, The γ -secretase structure is shown in cartoon representation (left) and surface view (right). Eleven N-linked glycans are displayed in stick. **b**, The γ -secretase structure is viewed perpendicular to the lipid membrane from the intracellular side. TM2 of PS1 is most flexible and shown in a semi-transparent fashion. The catalytic residues Asp257 and Asp385 are located on the convex side of the TM horseshoe. All structural figures were prepared using UCSF Chimera³⁵ or PyMol³⁶.

model building; nonetheless, we generated an atomic model for this TM on the basis of its sequence homology to the archaeal homologue PSH¹⁹. This is the first structure of human γ -secretase at a near-atomic resolution (Fig. 1 and Extended Data Fig. 3).

Nicastrin ECD, which constitutes the bulk of the extracellular region of human γ -secretase, contains 11 glycans and directly interacts with two ends of the TM horseshoe (Fig. 1a). The lone TM of nicastrin closely stacks against TM1/TM5/TM7 of APO-1 through predominantly hydrophobic interactions at the thick end. TM2 and TM4 of APO-1 associate with TM8 and TM9 of PS1 at the centre of the TM horseshoe, with the C-terminal three residues Phe465–Tyr466–Ile467 of PS1 inserting into a greasy central cavity on the extracellular side of APO-1 (Fig. 1a). The three TMs of PEN-2 bind to TM4 of PS1 at the thin end. Notably, this arrangement places the two catalytic residues, Asp257 and Asp385 of PS1, on the convex side of the TM horseshoe (Fig. 1b).

Atomic structure of presenilin

PS1 exhibits an extended organization, with empty spaces between a few adjacent TMs (Fig. 2a and Extended Data Fig. 4). The N-terminal 77 residues have no EM density and are presumably disordered, probably because of their intrinsic flexibility. Among the seven surface loops that connect neighbouring TMs, four exhibit clear and contiguous density. The extended sequences between TM6 and TM7 harbour the site of autocatalytic cleavage; but these sequences are hydrophilic and mostly disordered (Fig. 2b). The nine TMs exhibit a large variation in length, with TM9 containing 30 residues and TM7 only 18.

Among the two catalytic residues, Asp257 is located in the middle of TM6 slightly to the extracellular side, whereas Asp385 maps to the cytoplasmic side of TM7 (Fig. 2b). The distance between the C α atoms of Asp257 and Asp385 measures approximately 10.6 Å (Fig. 2c), considerably longer than that in an activated aspartate protease such as pepsin²². Importantly, however, these catalytic residues are placed




Figure 2 | Atomic structure of PS1. **a**, PS1 has a loosely organized structure and exhibits considerable flexibility. The cartoon representation of PS1 is rainbow-coloured. TM2 is visible only at low resolutions, and the density map contains no features for side chains. Nonetheless, an atomic model for TM2 was built on the basis of sequence and structural homology between PS1 and PSH^{19,21}. **b**, A membrane topology diagram of PS1. The two catalytic aspartate residues are coloured red. **c**, The two catalytic aspartate residues of PS1 are in near-perfect registry with those in PSH¹⁹. The PAL sequence motif implicated in substrate recognition is shown. **d**, PS1 and PSH share similar features at their active sites.

next to the PAL signature motif on TM9 that is thought to play a role in substrate recognition^{23–25} (Fig. 2c). We speculate that substrate binding may trigger alignment of these two aspartate residues and consequent catalysis.

Despite a relatively low sequence identity of 19% between PS1 and the archaeal intramembrane protease PSH, their overall structures are similar to each other^{19,21}. In particular, the catalytic residues are in nearly perfect registry between PS1 and PSH (Fig. 2c). The amino acids that surround the catalytic residues, including the PAL motif, are also highly conserved. Relative to Asp257, three residues of PS1 (Ile253, Tyr256, and Val261) are located on the same side of TM6; these residues are replaced by Leu158, Tyr161, and Val166, respectively, in PSH (Fig. 2d). Similarly, Gly382 and Phe388 of PS1 correspond to Gly219 and Met223 of PSH, respectively. Ile439 on TM9 of PS1 and Ile283 of PSH nearly coincide with each other (Fig. 2d). The observed structural conservation may underlie the finding that PSH exhibits similar cleavage preferences towards APP C99 as human γ -secretase²⁶.

Mutational hotspots in PS1 structure

The PS1 structure at a near-atomic resolution allows mapping and analysis of disease-derived mutations in PS1. On the basis of public information (<http://www.alzforum.org/mutations>), the 212

Alzheimer's-disease-derived missense mutations in PS1 affect 135 amino acids, of which 53 are targeted for two or more mutations. Presumably these residues affect the specific function of PS1 that is directly related to development of Alzheimer's disease. Among these 53 residues, 35 are identified in the TMs of our current PS1 structure, and the rest are located on surface elements that have no EM density. Notably, there are eight such residues in TM5, seven in TM6, but none in TM1. Together, these 35 amino acids account for a total of 101 Alzheimer's-disease-derived missense mutations.

Analysis of these 35 residues on the structure of PS1 led to identification of two mutational hotspots, each located at the inner core of a structural repeat (Fig. 3a). The first hotspot involves the inner core of TMs 2–5 (Fig. 3b). Among the 20 affected residues in these TMs, 18 have their side chains facing the inner core of TMs 2–5. Consequently, only one side of each TM helix is affected. For example, Leu219, Glu222, Leu226, Ser230, Met233, and Phe237 are placed on the same side of TM5 (Fig. 3b). Similarly, the second hotspot is located at the inner core of TMs 6–9, in the vicinity of the catalytic residues Asp257 and Asp385 (Fig. 3c).

To examine the functional consequence of disease-derived mutations, we generated ten γ -secretase variants, each containing a distinct missense mutation in PS1. Among these mutations, seven map to the two mutational hotspots. These γ -secretase variants were individually




Figure 3 | Alzheimer’s disease-derived mutations map to two hotspots in PS1. **a**, An overall view of the PS1 residues targeted for mutations in patients with Alzheimer’s disease. PS1 is viewed from the extracellular side. Mutated residues are coloured orange. **b**, Close-up views of the mutation-targeted residues in TMs 2–5. Most of these residues map to the centre of this four-TM bundle. **c**, Close-up views of the mutation-targeted residues in TMs 6–9. The two catalytic residues Asp257 and Asp385 are shown. **d**, FAD-derived mutations in PS1 have varying degrees of effect on the combined A β 40 and

Aβ42 cleavage activity of γ-secretase. Shown here are results of ten such γ-secretase mutants, each containing a specific mutation derived from FAD. The activity of WT γ-secretase is normalized to 1.0. **e**, FAD-derived mutations either suppressed the production of Aβ40 more than Aβ42 or increased the production of Aβ40 less than Aβ42. **f**, All but two FAD-derived mutations led to increased Aβ42/Aβ40 ratios. Two mutations F237I and V261F in PS1 abrogated Aβ40 cleavage altogether, disallowing calculation of the Aβ42/Aβ40 ratio. Each experiment was independently repeated three times. Error bars, s.d.




Figure 4 | Structural features of nicastrin. **a**, Two perpendicular views of nicastrin. The lid from the small lobe is highlighted in red. Surface glycans are shown. **b**, The lid hovers above a hydrophilic pocket in the large lobe. Two large glycans on Asn55 and Asn435 sandwich the lid and interact with surrounding residues. **c**, Glu333 and Tyr337 are surrounded by several charged and polar residues in the pocket. These structural features are consistent with the pocket being a binding site for substrate protein. **d**, Trp164 from the lid makes van der Waals contacts to Pro424, Phe448, and the aliphatic side chain of Gln420. **e**, Phe287 from the large lobe may serve as the hydrophobic pivot. Phe287 interacts with four hydrophobic residues from the small lobe.

purified to homogeneity and examined for their protease activities towards the APP C99 substrate in an *in vitro* cleavage assay. Compared with the wild-type (WT) γ -secretase, four mutations (I202F, F237I, L248R, and V261F) lead to severely compromised protease activity (Fig. 3d). Of these four mutations, F237I and V261F failed to generate any detectable amount of A β 40 (Fig. 3e), disallowing calculation of the A β 42/A β 40 ratio (Fig. 3f). In contrast, three mutations (I143V, F177L, and M233L) had relatively little impact on the total cleavage activity of γ -secretase (Fig. 3d), whereas the remaining three mutations (I213L, L226F, and L424V) actually increased the total cleavage activity (Fig. 3d). These observations strongly suggest a disconnection between the total protease activity of γ -secretase and the development of FAD (where these mutations were isolated). All eight mutations for which the A β 42/A β 40 ratio can be calculated led to increased A β 42/A β 40 ratios compared with WT γ -secretase (Fig. 3f). The generally increased ratios of A β 42 over A β 40 may suggest a causal relationship for development of FAD, but could also be explained by other possibilities.

Atomic structure of nicastrin

Similar to DpNCT²⁰, human nicastrin also contains a large lobe, a small lobe, and a lone TM (Fig. 4a and Extended Data Fig. 5). These two nicastrin structures can be aligned to each other with a root mean squared deviation of 2.2 Å. Of the 230 kDa molecular mass in mature γ -secretase, up to 70 kDa can be attributed to glycosylation of nicastrin²⁷. Among the 16 predicted N-linked glycosylation sites in nicastrin, at least 11 are glycosylated, as judged by the EM density. Because the extended glycans are flexible in solution, only a small portion close to each Asn residue was modelled (Fig. 4a).

Nicastrin is thought to rely on Glu333 and Tyr337 for substrate recruitment^{10–13}. In the structure of human nicastrin, Glu333 and Tyr337 are buried in a hydrophilic pocket that is covered by an extended surface loop known as the lid²⁰ (Fig. 4b). The lid, containing five aromatic residues, is sandwiched by two prominent glycans on Asn55 and Asn435 (Fig. 4b). Several charged and polar amino acids are located in the pocket, including four arginine residues: Arg281, Arg285, Arg429, and Arg432 (Fig. 4c). With the potential to mediate specific interactions such as hydrogen bonds, buried charged and/or

polar residues often serve a functional role. These structural features support the notion that this pocket is responsible for substrate recruitment, with these residues directly contributing to recognition.

The closed conformation of the lid is sustained by specific interactions between residues from the lid and residues from surrounding structural elements. For example, the indole ring of Trp164 makes several van der Waals contacts to the side chains of Pro424 and Phe448 and the aliphatic portion of Gln420 (Fig. 4d). Substrate binding requires opening of the lid and disruption of these interactions. Because the lid comes from the small lobe, a rotation of the large lobe around a central pivot—Phe287—is proposed to be both necessary and sufficient for lid opening²⁰. Such a rotation around Phe287 would be greatly facilitated by hydrophobic interactions, which are more adaptive to conformational changes than hydrogen bonds. Consistent with this analysis, Phe287 is nestled in a greasy pocket formed by Phe103, Leu171, Phe176, and Ile180 from the small lobe (Fig. 4e).

Inter-component interactions

Assembly of the four components into a functional human γ -secretase involves specific interactions mainly in the membrane-spanning region, resulting in the burial of approximately 3,794 Å² of otherwise membrane-exposed surface areas (Fig. 5a). Binding of the ECD onto the TMs of γ -secretase involves an additional buried surface area of 1,320 Å². The intra-membrane interactions comprise predominantly van der Waals contacts among hydrophobic amino acids, exemplified by the insertion of three residues at the C terminus of PS1 into a greasy pocket formed by TM2/3/4/6/7 of A β H-1 (Fig. 5b). Phe465 from PS1 contacts Leu72 and Phe205 in A β H-1, whereas Tyr466 from PS1 stacks against Leu163 and Ile167 in A β H-1. The aliphatic side chain of Ile467 from PS1 interacts with Leu46, Leu86, and Phe125 at the bottom of the cavity in A β H-1. The C terminus of PS1 is in close proximity to the highly conserved His171 and His197 in A β H-1, which were found to be important for A β H-1 binding to γ -secretase²⁸. Interestingly, a peptide comprising the C-terminal eight residues of PS1 was shown to inhibit A β production²⁹, presumably by weakening or disrupting normal assembly of γ -secretase.

On the thin end of the TM horseshoe, an α -helix and its preceding and ensuing loops in nicastrin stack against an extended loop at the




Figure 5 | Assembly interfaces among the four components of γ -secretase in the transmembrane region. **a**, An overall view of the packing interfaces in the transmembrane region. The three boxed interfaces are detailed in **b–d**. **b**, The C-terminal three residues Phe465–Tyr466–Ile467 of PS1 insert into a cavity formed by TMs in APH-1. **c**, Nicastrin interacts with PEN-2 through van der Waals contacts on a flat interface. **d**, PEN-2 binds to PS1 mostly through van der Waals contacts. In particular, Phe94 of PEN-2 is nestled in the greasy pocket of PS1, formed by Phe179, Leu182, Phe186, Val193, Tyr195, and Val198. **e**, Two phospholipids appear to stabilize the inter-component interfaces in γ -secretase. One lipid is bound at the interface between PS1 and APH-1, whereas the other is intercalated between the lone TM of nicastrin and TMs 1/4/5/7 of APH-1 (left). The aliphatic tails of the latter phospholipid may interact with several hydrophobic residues whereas its phosphate group likely hydrogen bonds to Arg115 and Gln116 (right).

C-terminal end of PEN-2 through multiple van der Waals contacts (Fig. 5c). At the nearby interface between PS1 and PEN-2, Phe94 from PEN-2 is nestled in a hydrophobic pocket formed by six residues in PS1 (Fig. 5d). Hydrophobic residues from TM3 of PEN-2, exemplified by Leu71, Ile75, and Phe78, also interact with residues in PS1. Notably, although PEN-2 has only three TMs and interacts with both PS1 and nicastrin, it actually contains three hydrophobic structural cores (Extended Data Fig. 6).

At 3.4 Å resolution, at least two phospholipid molecules were identified, each with two aliphatic tails linked to a small head group (Extended Data Fig. 2h). One lipid binds to the interface between PS1 and APH-1, making close contacts with residues in TM1/TM8 of PS1 and TM4 of APH-1 (Fig. 5e). The other lipid molecule is intercalated into the interface between APH-1 and the lone TM of nicastrin (Fig. 5e, right panel). The aliphatic tails of this phospholipid interact with several surrounding hydrophobic residues, whereas the phosphate group may directly hydrogen bond to the side chains of Arg115 and Gln116 from APH-1.

This lipid probably stabilizes the nicastrin–APH-1 interface within the membrane.

To corroborate the observed inter-component interfaces, we generated eight mutant γ -secretase complexes, each containing two specific cysteine mutations on two neighbouring components to facilitate formation of designed disulfide bonds. Three such γ -secretase mutants target the interface between APH-1 and nicastrin (Extended Data Fig. 7a). The distance between the C α atoms of Val147 on APH-1 and Ile40 on nicastrin is 4.1 Å, ideal for formation of a disulfide bond if these two residues were replaced by cysteine. Indeed, in the absence of the reducing agent DTT, APH-1 V147C was crosslinked to nicastrin I40C within the mutant γ -secretase, resulting in a high molecular mass complex on denaturing SDS–polyacrylamide gel electrophoresis (SDS–PAGE) (Extended Data Fig. 7a, lane 1). This crosslinked complex disappeared in the presence of DTT (lane 2) and was absent for the WT γ -secretase regardless of DTT (lanes 7 and 8). Similarly, two additional γ -secretase mutants, one containing APH-1 V146C and nicastrin A664C and the other containing APH-1 A4C and nicastrin L673C, allowed specific crosslinking only in the absence of DTT (lanes 3–6). The same strategy was successfully applied to verify the specific interactions at the interfaces between PS1 and PEN-2 (Extended Data Fig. 7b), between PEN-2 and nicastrin (Extended Data Fig. 7c), and between APH-1 and PS1 (Extended Data Fig. 7d).

Discussion

In this paper, we report the cryo-EM structure of human γ -secretase at an overall resolution of 3.4 Å. The qualitative improvement in resolution over earlier studies allowed us to derive an atomic model for all four components of γ -secretase. Except nicastrin^{18,20}, such atomic models are reported here for the first time.

The available structural evidence supports the notion that Glu333 and Tyr337 of nicastrin, both buried in a hydrophilic pocket, may play a key role in substrate recruitment¹². Under this scenario, displacement of the lid, caused by a movement of the large lobe relative to the small lobe, is required before substrate binding²⁰. This proposal is supported by the unique pattern of contacts between the large and small lobes and by the conversed interactions around the central pivot Phe287 in nicastrin (Fig. 4). In substrate-free γ -secretase, the lid is positioned right above the concave side of the TM horseshoe whereas the active site is located on the convex side, resulting in a relatively long distance between the putative binding pocket for the N terminus of the substrate and the site of cleavage by PS1 (Extended Data Fig. 8). This distance may shorten, however, in response to conformational changes within γ -secretase. In particular, rotation of the large lobe relative to the small lobe, induced by substrate binding, may also re-orient the substrate for cleavage, perhaps by aligning the pocket in nicastrin and the active site in PS1. Dynamic conformations of γ -secretase have been observed³⁰, particularly in the ECD region³¹.

TM6 of PS1, which harbours the catalytic residue Asp257, exhibits relatively poor EM density, whereas TM2 is visible only at low resolutions. These structural observations are consistent with the notion that binding by inhibitors or modulators may induce pronounced conformational changes^{30–32}. In particular, substrate binding may trigger a conformational change, rendering the active site suitable for catalysis³². The plasticity observed in our structure may expedite the conformational changes that are needed to bring the two catalytic aspartates within hydrogen bonding distance of each other, and may play a role in the relaxed substrate specificity of the complex. Activation of the active site also depends on the binding of PEN-2, which was observed to have an allosteric effect on TM6 (ref. 33).

The prevailing β -amyloid hypothesis suggests an increased ratio of A β 42 over A β 40 as the major culprit for the accumulation of β -amyloid plaque and consequent development of Alzheimer's disease^{2,3}. Our finding that FAD-derived mutations have varying degrees of effect on the protease activity of γ -secretase argues against a causal

relationship between the total protease activity of γ -secretase and the development of FAD. Our finding is consistent with the poor clinical performance of γ -secretase-inhibiting drugs³⁴. Notably, the generally increased ratios of A β 42 over A β 40 only suggest, but fail to prove, a direct causal relationship for the development of FAD. For example, the assumption that γ -secretase has evolved to optimize production of A β 40 over all other peptides would naturally lead to the prediction that any mutation, disease-causing or not, will increase the A β 42/A β 40 ratio.

Intriguingly, the mutations causing Alzheimer's disease map to two hotspots, each located at the inner core of a four-TM bundle (Fig. 3). The mutations on TMs 6–9 affect residues in the immediate neighbourhood of the catalytic residues Asp257 and Asp385 and thus may cripple the protease activity of γ -secretase. The biochemical data seem to support this analysis (Fig. 3d). However, the mutations on TMs 2–5 defy such rationale: they either abrogate or increase the protease activity of γ -secretase. The generally inward-facing nature of the mutation-targeted residues on TMs 2–5 (Fig. 3b) may suggest a transport function of some sort, or a binding site for another transmembrane protein. These tantalizing clues await experimental examination.

Our structure represents a milestone for the rapidly emerging technique of high-resolution cryo-EM structure determination. Because the signal-to-noise ratio in cryo-EM images correlates with the size of the particles, determining the relative orientations of individual particles becomes a limitation for small complexes. With a protein mass of ~170 kDa, γ -secretase is much smaller than any other atomic-resolution cryo-EM structure determined so far, while its lack of symmetry further complicates structure determination. Whereas glycosylation adds another 30–70 kDa of mass to the complex, only a small proportion of the sugar moieties are ordered and their effect on alignment is limited. Although glycosylation may inhibit crystal growth, it did not seem to hamper cryo-EM structure determination. On the contrary, glycosylation may stabilize the protein, possibly by protecting the protein from the denaturing air–water interface of the thin cryo-EM sample.

Online Content Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

Received 10 May; accepted 6 July 2015.

Published online 17 August 2015.

1. Alzheimer, A. About a peculiar disease of the cerebral cortex. *Zbl. Nervenheilk. Psychiatrie* **30**, 177–179 (1907).
2. Hardy, J. A. & Higgins, G. A. Alzheimer's disease: the amyloid cascade hypothesis. *Science* **256**, 184–185 (1992).
3. De Strooper, B., Iwatsubo, T. & Wolfe, M. S. Presenilins and γ -secretase: structure, function, and role in Alzheimer disease. *Cold Spring Harb. Perspect. Med.* **2**, a006304 (2012).
4. Goate, A. & Hardy, J. Twenty years of Alzheimer's disease-causing mutations. *J. Neurochem.* **120** (suppl. 1), 3–8 (2012).
5. Tanzi, R. E. & Bertram, L. Twenty years of the Alzheimer's disease amyloid hypothesis: a genetic perspective. *Cell* **120**, 545–555 (2005).
6. Kimberly, W. T. et al. γ -Secretase is a membrane protein complex comprised of presenilin, nicastrin, aph-1, and pen-2. *Proc. Natl Acad. Sci. USA* **100**, 6382–6387 (2003).
7. De Strooper, B. et al. Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature* **391**, 387–390 (1998).
8. Wolfe, M. S. et al. Two transmembrane aspartates in presenilin-1 required for presenilin endoproteolysis and γ -secretase activity. *Nature* **398**, 513–517 (1999).
9. Thinakaran, G. et al. Endoproteolysis of presenilin 1 and accumulation of processed derivatives in vivo. *Neuron* **17**, 181–190 (1996).
10. Yu, G. et al. Nicastrin modulates presenilin-mediated notch/glp-1 signal transduction and β APP processing. *Nature* **407**, 48–54 (2000).
11. Shah, S. et al. Nicastrin functions as a γ -secretase-substrate receptor. *Cell* **122**, 435–447 (2005).
12. Dries, D. R. et al. Glu-333 of nicastrin directly participates in γ -secretase activity. *J. Biol. Chem.* **284**, 29714–29724 (2009).
13. Goo, J. S. et al. Nicastrin overexpression in transgenic mice induces aberrant behavior and APP processing. *Mol. Neurobiol.* **48**, 232–243 (2013).

14. Francis, R. et al. *aph-1* and *pen-2* are required for Notch pathway signaling, γ -secretase cleavage of β APP, and presenilin protein accumulation. *Dev. Cell* **3**, 85–97 (2002).
15. Takasugi, N. et al. The role of presenilin cofactors in the γ -secretase complex. *Nature* **422**, 438–441 (2003).
16. Goutte, C., Tsunozaki, M., Hale, V. A. & Priess, J. R. APH-1 is a multipass membrane protein essential for the Notch signaling pathway in *Caenorhabditis elegans* embryos. *Proc. Natl Acad. Sci. USA* **99**, 775–779 (2002).
17. Gu, Y. et al. APH-1 interacts with mature and immature forms of presenilins and nicastrin and may play a role in maturation of presenilin-nicastrin complexes. *J. Biol. Chem.* **278**, 7374–7380 (2003).
18. Lu, P. et al. Three-dimensional structure of human γ -secretase. *Nature* **512**, 166–170 (2014).
19. Li, X. et al. Structure of a presenilin family intramembrane aspartate protease. *Nature* **493**, 56–61 (2013).
20. Xie, T. et al. Crystal structure of the γ -secretase component nicastrin. *Proc. Natl Acad. Sci. USA* **111**, 13349–13354 (2014).
21. Sun, L. et al. Structural basis of human γ -secretase assembly. *Proc. Natl Acad. Sci. USA* **112**, 6003–6008 (2015).
22. Cooper, J. B., Khan, G., Taylor, G., Tickle, I. J. & Blundell, T. L. X-ray analyses of aspartic proteinases. II. Three-dimensional structure of the hexagonal crystal form of porcine pepsin at 2.3 Å resolution. *J. Mol. Biol.* **214**, 199–222 (1990).
23. Wang, J., Brunkan, A. L., Hecimovic, S., Walker, E. & Goate, A. Conserved “PAL” sequence in presenilins is essential for γ -secretase activity, but not required for formation or stabilization of γ -secretase complexes. *Neurobiol. Dis.* **15**, 654–666 (2004).
24. Sato, C., Takagi, S., Tomita, T. & Iwatsubo, T. The C-terminal PAL motif and transmembrane domain 9 of presenilin 1 are involved in the formation of the catalytic pore of the γ -secretase. *J. Neurosci.* **28**, 6264–6271 (2008).
25. Wang, J. et al. C-terminal PAL motif of presenilin and presenilin homologues required for normal active site conformation. *J. Neurochem.* **96**, 218–227 (2006).
26. Dang, S. et al. Cleavage of amyloid precursor protein by an archaeal presenilin homologue PSH. *Proc. Natl Acad. Sci. USA* **112**, 3344–3349 (2015).
27. Schedin-Weiss, S., Winblad, B. & Tjernberg, L. O. The role of protein glycosylation in Alzheimer disease. *FEBS J.* **281**, 46–62 (2014).
28. Pardossi-Piquard, R. et al. APH1 polar transmembrane residues regulate the assembly and activity of presenilin complexes. *J. Biol. Chem.* **284**, 16298–16307 (2009).
29. Esselens, C. et al. Peptides based on the presenilin-APP binding domain inhibit APP processing and A β production through interfering with the APP transmembrane domain. *FASEB J.* **26**, 3765–3778 (2012).
30. Elad, N. et al. The dynamic conformational landscape of γ -secretase. *J. Cell Sci.* **128**, 589–598 (2015).
31. Li, Y. et al. Structural interactions between inhibitor and substrate docking sites give insight into mechanisms of human PS1 complexes. *Structure* **22**, 125–135 (2014).
32. Takagi-Niidome, S. et al. Cooperative roles of hydrophilic loop 1 and the C-terminus of presenilin 1 in the substrate-gating mechanism of γ -secretase. *J. Neurosci.* **35**, 2646–2656 (2015).
33. Takeo, K., Watanabe, N., Tomita, T. & Iwatsubo, T. Contribution of the γ -secretase subunits to the formation of catalytic pore of presenilin 1 protein. *J. Biol. Chem.* **287**, 25834–25843 (2012).
34. De Strooper, B. Lessons from a failed γ -secretase Alzheimer trial. *Cell* **159**, 721–726 (2014).
35. Pettersen, E. F. et al. UCSF Chimera—a visualization system for exploratory research and analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).
36. DeLano, W. L. The PyMOL Molecular Graphics System (Schrödinger, LLC, 2002).

Acknowledgements We thank S. Chen and C. Savva for support with electron microscopy, and J. Grimmer and T. Darling for support with high-performance computing. This work was supported by funds from the Ministry of Science and Technology (2014ZX09507003006 to Y.S.), the National Natural Science Foundation of China (31130002 and 31321062 to Y.S.), a European Union Marie Curie Fellowship (to X.C.B.), and the UK Medical Research Council (MC_UP_A025_1013, to S.H.W.S.).

Author Contributions Y.S. initiated and supervised the project. G.Y., P.L., D.M., L.S., and R.Z. prepared the sample and pre-screened samples in various detergents on F20. X.B. prepared grids and collected cryo-EM data. X.B. and S.S. calculated the cryo-EM map. C.Y. built and refined the atomic model. X.B. independently built and refined the atomic model. Y.S., L.S., G.Y., and R.Z. designed and analysed the mutational and biochemical characterizations. L.S., G.Y., and R.Z. performed the biochemical assays. All authors contributed to analysis of the structure. X.B., C.Y., S.S. and Y.S. contributed to manuscript preparation.

Author Information The atomic coordinates have been deposited in the Protein Data Bank under accession number 5A63, and the EM maps have been deposited in the Electron Microscopy Data Bank under accession code EMD-3061. Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Correspondence and requests for materials should be addressed to X.B. (xcbai@mrc-lmb.cam.ac.uk), S.S. (scheres@mrc-lmb.cam.ac.uk), or Y.S. (shi-lab@tsinghua.edu.cn).

METHODS

No statistical methods were used to predetermine sample size. The experiments were not randomized. The investigators were not blinded to allocation during experiments and outcome assessment.

Protein preparation. To improve protein production, we co-expressed the four components of γ -secretase, each in its own pMLink plasmid¹⁸. This strategy allowed both convenient manipulation of individual components and optimization of protein expression by altering the relative molar ratios of the four plasmids. Specific mutations in PS1, PEN-2, APH-1, or nicastrin were generated only in the corresponding pMLink vector using a standard PCR-based approach. PEN-2 has an amino-terminal Flag tag, APH-1 contains a carboxyl-terminal haemagglutinin (HA) tag, and nicastrin is tagged with a V5-His₆ sequence at the carboxyl terminus. The ten γ -secretase variants, each carrying a FAD-derived mutation on PS1, and the eight γ -secretase complexes, each carrying two introduced cysteine mutations, were overexpressed similarly as the WT γ -secretase¹⁸. Purification of both WT and mutant human γ -secretases followed published protocol¹⁸.

Electron microscopy. Aliquots of 3 μ l of purified γ -secretase in amphipols at a concentration of $\sim 4 \mu$ M were placed on glow-discharged holey carbon grids (Quantifoil Au R1.2/1.3, 300 mesh), and flash frozen in liquid ethane using an FEI Vitrobot. Zero-energy-loss images were recorded manually on an FEI Titan Krios electron microscope at 300 kV, using a slit width of 20 eV on a GIF-Quantum energy filter. A Gatan K2-Summit detector was used in super-resolution counting mode at a calibrated magnification of $\times 35,714$ (yielding a pixel size of 1.4 Å), and a dose rate of ~ 2.5 electrons per square ångström per second (approximately 5 electrons per pixel per second). Exposures of 16 s were dose-fractionated into 20 movie frames. Defocus values in the final data set ranged from 0.7 to 3.2 μ m.

Image processing. We used MOTIONCORR³⁷ for whole-frame motion correction, CTFFIND3 (ref. 38) for estimation of the contrast transfer function parameters, and RELION-1.3 (ref. 39) for all other image processing steps. Templates for reference-based particle picking were obtained from 2D class averages that were calculated from a manually picked subset of the micrographs. Using low-pass filtered templates to 20 Å to limit reference bias, 1.8 million particles were picked automatically from a total of 2,925 micrographs. Because the picking procedure is prone to false positives⁴⁰, we used reference-free 2D class averaging and an initial run of 3D classification to select 412,272 particles for a first 3D refinement. After per-particle motion correction and radiation-damage weighting (particle polishing⁴¹), these particles gave a reconstruction with a resolution of 3.5 Å. In a subsequent 3D classification, we applied a mask around the transmembrane domain and did not perform any alignments. This yielded a subset of 159,549 particles, for which a reconstruction with improved density in the trans-membrane domain was obtained. The resolution of this final map was 3.4 Å. Reported resolutions are based on the gold-standard FSC = 0.143 criterion³⁹, and FSC curves were corrected for the effects of a soft mask on the FSC curve using high-resolution noise substitution⁴². All 3D classifications and refinements were started from a 40 Å low-pass filtered initial model, the first of which was made from our previous 4.5 Å map. Before visualization, all density maps were corrected for the modulation transfer function of the detector, and then sharpened by applying a negative B-factor that was estimated using automated procedures⁴³. Local resolution variations were estimated using ResMap⁴⁴.

Atomic modelling. The first atomic model was built from an intermediate, 3.9 Å EM map which was of sufficient quality to allow side-chain assignment for all four components of γ -secretase. The initial models used for nicastrin ECD and presenilin 1 were generated from the crystal structures of DpNCT²⁰ (Protein Data Bank accession number 4R12) and PSH¹⁹ (Protein Data Bank accession number 4HYG) by CHAINSAW⁴⁵. The two generated structures were first refined in real-space by PHENIX⁴⁶ with secondary structure and geometry restraints. APH-1 and PEN-2 were built *de novo* from a poly-Ala model. Then the entire atomic model was manually improved in COOT⁴⁷. Sequence assignment was guided mainly by bulky residues such as Phe, Tyr, Trp and Arg. Unique patterns of sequences were exploited for validation of residue assignment. Eleven glycosylation sites with obvious sugar densities and five pairs of disulfide bonds in the nicastrin ECD domain also facilitated sequence assignment. This initial model


was largely confirmed and further improved by the 3.4 Å resolution EM map, with minor adjustment to a few residues and identification of two additional glycosylation sites and two lipid molecules. This model was refined in real-space by PHENIX⁴⁶.

An independent effort of atomic model building used stereochemical refinement tools in Coot⁴⁷ and REFMAC⁴⁸ that were originally designed for X-ray crystallography, but later adapted for cryo-EM⁴⁹. The structures of DpNCT²⁰ (Protein Data Bank accession number 4R12) and PSH¹⁹ (Protein Data Bank accession number 4HYC) were used as a guide, while APH-1 and PEN-2 were built *de novo* in Coot⁴⁷, starting from idealized α -helices. This model was refined in REFMAC⁴⁸, with secondary structure restraints generated by ProSMART⁵⁰. Superposition of the two independently built models revealed that both models were in excellent agreement. Refinement of the final, consensus model was performed in REFMAC, and overfitting of the atomic model was monitored by refining the model in one of the two independent maps from the gold-standard refinement approach, and testing this model against the other map⁵¹. At no point in our procedures did we use the atomic model to modify the cryo-EM map.

γ -Secretase activity assay. The assay was performed as described in the AlphaLISA Kit (PerkinElmer). Briefly, 2 μ l reaction products were incubated with 8 μ l AlphaLISA AB1-40/42 Acceptor beads at 23 °C for 1 h. After another incubation for 30 min with 10 μ l AlphaLISA AB1-40/42 Donor beads in the dark at 23 °C, the samples were read using Envision-Alpha Reader (PerkinElmer). The readings were expressed in arbitrary units.


Crosslinking assay. For the crosslinking assay, copper sulfate was added to the purified protein at a final concentration of 1 mM, and the sample was incubated at 4 °C for 1 h. Then, the protein was applied to gel filtration (Superose-6, GE Healthcare). The peak fractions were divided into two aliquots. One aliquot was treated with the reducing agent DL-dithiothreitol (DTT), at a final concentration of 10 mM and incubated at room temperature for 30 min. Samples from the two aliquots, treated with or without DTT, were analysed by western blotting using monoclonal antibodies against the HA tag on APH-1 or Flag tag on PEN-2. The antibodies were purchased from Beijing ComWin Biotech.

37. Li, X. *et al.* Electron counting and beam-induced motion correction enable near-atomic-resolution single-particle cryo-EM. *Nature Methods* **10**, 584–590 (2013).
38. Mindell, J. A. & Grigorieff, N. Accurate determination of local defocus and specimen tilt in electron microscopy. *J. Struct. Biol.* **142**, 334–347 (2003).
39. Scheres, S. H. RELION: implementation of a Bayesian approach to cryo-EM structure determination. *J. Struct. Biol.* **180**, 519–530 (2012).
40. Scheres, S. H. Semi-automated selection of cryo-EM particles in RELION-1.3. *J. Struct. Biol.* **189**, 114–122 (2015).
41. Scheres, S. H. Beam-induced motion correction for sub-megadalton cryo-EM particles. *eLife* **3**, e03665 (2014).
42. Chen, S. *et al.* High-resolution noise substitution to measure overfitting and validate resolution in 3D structure determination by single particle electron cryomicroscopy. *Ultramicroscopy* **135**, 24–35 (2013).
43. Rosenthal, P. B. & Henderson, R. Optimal determination of particle orientation, absolute hand, and contrast loss in single-particle electron cryomicroscopy. *J. Mol. Biol.* **333**, 721–745 (2003).
44. Kucukelbir, A., Sigworth, F. J. & Tagare, H. D. Quantifying the local resolution of cryo-EM density maps. *Nature Methods* **11**, 63–65 (2014).
45. Stein, N. CHAINSAW: a program for mutating pdb files used as templates in molecular replacement. *J. Appl. Cryst.* **41**, 641–643 (2008).
46. Adams, P. D. *et al.* PHENIX: building new software for automated crystallographic structure determination. *Acta Crystallogr. D* **58**, 1948–1954 (2002).
47. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr. D* **60**, 2126–2132 (2004).
48. Murshudov, G. N., Vagin, A. A. & Dodson, E. J. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr. D* **53**, 240–255 (1997).
49. Brown, A. *et al.* Tools for macromolecular model building and refinement into electron cryo-microscopy reconstructions. *Acta Crystallogr. D* **71**, 136–153 (2015).
50. Nicholls, R. A., Fischer, M., McNicholas, S. & Murshudov, G. N. Conformation-independent structural comparison of macromolecules with ProSMART. *Acta Crystallogr. D* **70**, 2487–2499 (2014).
51. Amunts, A. *et al.* Structure of the yeast mitochondrial large ribosomal subunit. *Science* **343**, 1485–1489 (2014).
52. Crystal, A. S. *et al.* Membrane topology of γ -secretase component PEN-2. *J. Biol. Chem.* **278**, 20117–20123 (2003).




Extended Data Figure 1 | Cryo-EM, single-particle analysis of human γ -secretase. **a**, Representative raw particles from an original micrograph. **b**, Representative reference-free 2D class averages of the γ -secretase particles. Two classes identified by a red rectangle box (lower right corner) may contain some density for the extended cytosolic loop sequences between TM6 and TM7 of PS1, which are disordered in the final maps. **c**, Resolution estimation of the EM structure. The overall resolution is calculated to be 3.4 Å on the basis of gold-standard FSC curve³⁹. **d**, Colour-coded resolution variations in the

γ -secretase structure as estimated by ResMap⁴⁴. **e**, FSC curves of the final, Refmac-refined model versus the map it was refined against (in black); of a model refined in the first of the two independent maps used for the gold-standard FSC versus that same map (in red); and of a model refined in the first of the two independent maps versus the second independent map (in green). The small difference between the red and green curves indicates that the refinement of the atomic coordinates did not suffer from severe overfitting.



Extended Data Figure 2 | An atomic model of human γ -secretase. **a**, The γ -secretase structure is viewed parallel to the lipid membrane. Shown here is EM density for the entire γ -secretase complex. EM density is coloured blue for PS1, yellow for PEN-2, magenta for APH-1, and green for nicastrin. **b**, The density map for TM2 of PS1. Among the 20 TMs, TM2 of PS1 shows the highest degree of flexibility and only becomes visible at as rod-shaped density in a 7 Å low-pass filtered map. At this resolution, another rod-shaped density is visible next to TM2 and remains unaccounted for. **c**, EM density map and the


atomic model are shown for all seven TMs of APH-1. Two to three bulky residues are indicated for each TM. **d**, EM density map and the atomic model are shown for seven TMs of human PS1. TM6 exhibits relatively poor EM density, probably because of its intrinsic flexibility. **e**, EM density map and the atomic model are shown for the three TMs of PEN-2. **f**, EM density map and the atomic model are shown for the only TM and select regions of nicastrin. **g**, EM density map and the atomic model for three representative glycans. **h**, EM density map and putative assignment for two lipid molecules.




Extended Data Figure 3 | Overall structure of human γ -secretase.

a, Structure of human γ -secretase is shown in cartoon representation (top) and surface view (bottom) in four successively perpendicular views. The γ -secretase structure is viewed parallel to the lipid membrane. The colouring scheme is

the same as in Fig. 1. Two lipid molecules are shown. Eleven glycosylated Asn residues and their glycans are displayed in stick. **b**, The γ -secretase structure is represented by electrostatic surface potential.




Extended Data Figure 4 | Electrostatic surface potential of PS1. PS1 exhibits a loosely folded structure, with several large cavities and empty spaces between adjacent TMs.




Extended Data Figure 5 | Structural comparison between human nicastrin and nicastrin from *D. purpureum* (DpNCT). Individual structures of human nicastrin and DpNCT²⁰ are shown in the left and middle panels,

respectively. The overlay is shown in the right panel, with a root mean squared deviation of 2.2 Å. Two perpendicular views for each structure are displayed here.




Extended Data Figure 6 | PEN-2 contains three small hydrophobic cores in its three TMs. Unlike previous prediction^{14,52}, PEN-2 contains three, not two, TMs. PEN-2 contains a small hydrophobic core in the extracellular side

and two in the transmembrane region. These three regions are boxed and shown in close-up views.



Extended Data Figure 7 | Results of crosslinking experiments corroborate the atomic model of γ -secretase. **a**, Crosslinking results for the interface between APh-1 and nicastrin (NCT). Three mutant γ -secretase complexes were examined: APh-1-V147C/NCT-I40C, APh-1-V146C/NCT-A664C, and APh-1-A4C/NCT-L673C. Shown in the upper panel is an SDS-PAGE gel blotted by a monoclonal antibody against the HA tag on APh-1. Only in the absence of DTT did crosslinking lead to high molecular mass complexes for the mutant γ -secretase, but not for the WT γ -secretase. The two bands probably represent APh-1 crosslinked to mature nicastrin (mNCT) and immature nicastrin (iNCT). The structural basis is shown in the lower panel. The

distances between the C α atoms of the two residues targeted for cysteine mutation range from 4.1 to 6.1 Å, which would facilitate convenient crosslinking reactions. **b**, Crosslinking results for the interface between PS1 and PEN-2. The mutant γ -secretase contains PEN-2-P97C and PS1-N190C. Shown in the upper panel is an SDS-PAGE gel blotted by a monoclonal antibody against the Flag tag on PEN-2. **c**, Crosslinking results for the interface between PEN-2 and nicastrin. Two γ -secretase mutants were examined: PEN-2-T100C/NCT-V224C and PEN-2-L98C/NCT-H222C. **d**, Crosslinking results for the interface between APh-1 and PS1. Two γ -secretase mutants were examined: APh-1-T204C/PS1-F465C and APh-1-A76C/PS1-I467C.



Extended Data Figure 8 | Implication on substrate access to γ -secretase. Structure of γ -secretase is displayed in three relevant views: left, electrostatic surface potential from the convex side of γ -secretase; middle, overall structure,

with key features labelled; right, suggested putative path for substrate access to the active site of γ -secretase.

Extended Data Table 1 | Refinement and model statistics

Data Collection	
Particles	159,549
Pixel size (Å)	1.4
Defocus range (μm)	0.7-3.2
Voltage (kV)	300
Electron dose (e-/Å ⁻²)	40
Model composition	
Non-hydrogen atoms	10,016
Protein residues	1,251
Phospholipids (PC)	2
Sugars	15
Refinement	
Resolution (Å)	3.4
Map sharpening B-factor (Å ²)	-109
Fourier Shell Correlation	0.8503
Rfactor	0.3494
Rms deviations	
Bonds (Å)	0.0077
Angles (°)	1.4224
Validation (proteins)	
Molprobit score	2.65
Clashscore, all atoms	8.22
Good rotamers (%)	93.5%
Ramachandran plot	
Favored (%)	88.5
Allowed (%)	8.6
Outliers (%)	2.9